

# THE CLAIMS

## We claim:


1. An amphiphilic drug-oligomer conjugate comprising a therapeutic compound conjugated to an oligomer, wherein the oligomer comprises a lipophilic moiety coupled to a hydrophilic moiety.
2. The amphiphilic drug-oligomer conjugate of claim 1 further characterized in that said conjugate exhibits activity without cleavage of the therapeutic compound from the oligomer.
3. The amphiphilic drug-oligomer conjugate of claim 1 further characterized in that said therapeutic compound does not exhibit activity without cleavage of the therapeutic compound from the oligomer.
4. The amphiphilic drug-oligomer conjugate of claim 1 wherein the therapeutic compound is a peptide or protein.
5. The amphiphilic drug-oligomer conjugate of claim 1 wherein the therapeutic compound is a peptide and the peptide is selected from the group consisting of: enkephalin, adrenocorticotrophic hormone, adenosine deaminase ribonuclease, alkaline phosphatase, angiotensin, antibodies, arginase, arginine deaminase, asparaginase, caerulein, calcitonin, chemotrypsin, cholecystokinin, clotting factors, dynorphins, endorphins, endorphins, enkephalins, enkephalins, erythropoietin, gastrin-releasing peptide, glucagon, hemoglobin, hypothalamic releasing factors, interferon, katalcalcin, motilin, neuropeptide Y, neurotensin, non-naturally occurring opioids, oxytosin, papain, parathyroid hormone, peptides prolactin, soluble CD-4, somatomedin, somatostatin, somatostatin, somatotropin, superoxide dismutase, thyroid stimulating hormone, tissue plasminogen activator, trypsin, vasopressin and analogues and active fragments thereof.

Sub  
aa1


- Sub  
aa1  
cont
6. The amphiphilic drug-oligomer conjugate of claim 1 wherein the therapeutic compound is an Opioid receptor agonist, antagonist or partial agonist/partial antagonist.
- 5 7. The amphiphilic drug-oligomer conjugate of claim 1 wherein the therapeutic compound is [met<sup>51</sup>]enkephalin.
8. The amphiphilic drug-oligomer conjugate of claim 1 wherein the lipophilic moiety is coupled to the hydrophilic moiety by a hydrolyzable bond.
9. The amphiphilic drug-oligomer conjugate of claim 1 wherein the lipophilic moiety is coupled to the hydrophilic moiety by a non-hydrolyzable bond.
- 10 10. The The amphiphilic drug-oligomer conjugate of claim 1 wherein the lipophilic moiety is coupled to the hydrophilic moiety by a bond selected from the group consisting of: amide bond, carbamate bond, carbonate bond and ester bond.
- Sub  
aa2
11. The amphiphilic drug-oligomer conjugate of claim 1 wherein the oligomer moiety is coupled to the drug moiety by a bond selected from the group consisting of amide bond, carbamate bond, carbonate, and ester.
- 15 12. The amphiphilic drug-oligomer conjugate of claim 1 wherein the lipophilic moiety is selected from the group consisting of fatty acids, C<sub>1-26</sub>alkyls, cholesterol, and adamantane.
13. The amphiphilic drug-oligomer conjugate of claim 1 wherein the hydrophilic moiety is selected from the group consisting of sugars or PEG<sub>1-7</sub>.
- 20 14. The amphiphilic drug-oligomer conjugate of claim 1 wherein the hydrophilic moiety comprises a sugar and the sugar is selected from the group consisting of: amino sugars and non-amino sugars.
- Sub  
aa3
15. The amphiphilic drug-oligomer conjugate of claim 1 wherein the oligomer is selected from the group consisting of:
- 25



wherein  $n=3$  to 25 and  $m=1$  to 6;



wherein  $n=3$  to 25 and  $m=1$  to 7;




wherein  $n=3$  to 25,  $m=1$  to 7 and  $X=\text{O}$  or  $\text{N}$ ;



wherein  $m=0$  to 5 and  $\text{R}=\text{cholesterol}$  or  $\text{adamantane}$ ; or



wherein  $m=0$  to 5;



wherein  $m=0$  to 7;



wherein  $m=1$  to 7 and  $X=\text{N}$  or  $\text{O}$ .


- 15 16. The method of claim 1 wherein the hydrophilic moiety comprises a sugar and the sugar is selected from the group consisting of amino sugars and non-amino sugars.

17. The method of claim 1 wherein the hydrophilic moiety comprises a monosaccharide.
18. An amphiphilic drug-oligomer conjugate of claim 1 wherein the oligomer is selected from the group consisting of:



where  $m=1$  to 7;


Sub  
aa3  
cont




where  $m=1$  to 7;




10 where  $m=1$  to 6;




where  $m=1$  to 6;



where  $m=1$  to 6;




where  $m=1$  to 7;



where  $m=1$  to 7;



where  $m=1$  to 6;




where  $m=1$  to 6;




5

where  $m=1$  to 6;



where  $m=1$  to 6;




where  $m=1$  to 7;


10



where  $m=1$  to 6;



where  $m=1$  to 6;




15

where  $m=1$  to 6;



where  $m=1$  to 7;



Sub  
add  
cont

where  $m=1$  to 7;



where  $m=1$  to 6;






where  $m=1$  to 6; and




where  $m=1$  to 6.

19. The amphiphilic drug-oligomer conjugate of claim 1 wherein the therapeutic compound is a peptide having an added N-terminal residue selected from the group consisting of proline or alanine.

20. An amphiphilic oligomer-enkephalin conjugate selected from the group consisting of:




5



10

Sub  
day  
over





15

20

and


25



30

21. An amphiphilic oligomer-enkephalin conjugate wherein the oligomer is comprised of a lipophile and a hydrophile and the lipophile is coupled to the hydrophile by a hydrolyzable bond, said conjugate being selected from the group consisting of:


35




40



45






5

10



22. An amphiphilic oligomer-enkephalin conjugate wherein the oligomer is comprised of a lipophile and a hydrophile and the lipophile is coupled to the hydrophile by a non-hydrolyzable bond, said conjugate being selected from the group consisting of:

15




20


25



30



35



40

23. A method for activating a receptor comprising bringing said receptor into contact with an amphiphilic drug-oligomer conjugate comprising a therapeutic compound conjugated to an oligomer, wherein the oligomer comprises a lipophilic moiety coupled with to a hydrophilic moiety.



- Sub  
AA5
- 5
24. The method of claim 18 further characterized in that said conjugate exhibits activity in the without cleavage of the therapeutic compound from the oligomer.
25. The method of claim 18 wherein the receptor is a G-protein coupled receptor.
26. The method of claim 18 wherein the receptor is an Opioid receptor.
27. The method of claim 18 wherein the receptor is a Opioid receptor; selected from the group consisting of  $\delta$ ,  $\mu$ , and  $\kappa$ .
28. The method of claim 18 wherein the hydrophilic moiety is selected from the group consisting of sugar and PEG<sub>1-7</sub>.
29. The method of claim 18 wherein the hydrophilic moiety is selected from the group consisting of fatty acid, alkyl 1-26, cholesterol and adamantane.
30. The method of claim 18 wherein the therapeutic compound is a peptide having an added N-terminal residue selected from the group consisting of proline, and alanine.
31. The method of claim 18 wherein the therapeutic compound is a peptide or protein.
32. The method of claim 18 wherein the therapeutic compound is a peptide and the peptide is selected from the group consisting of: enkephalin, adrenocorticotrophic hormone, adenosine deaminase ribonuclease, alkaline phosphatase, angiotensin, antibodies, arginase, arginine deaminease, asparaginase, caerulein, calcitonin, chemotrypsin, cholecystokinin, clotting factors, dynorphins, endorphins, endorphins, enkephalins, enkephalins, erythropoietin, gastrin-releasing peptide, glucagon, hemoglobin, hypothalamic releasing factors, interferon, katalcalcin, motilin, neuropeptide Y, neurotensin, non-naturally occurring opioids, oxytosin, papain, parathyroid hormone, peptides prolactin, soluble CD-4, somatomedin, somatostatin, somatostatin, somatotropin, superoxide dismutase, thyroid stimulating hormone, tissue plasminogen

activator, trypsin, vasopressin, and analogues and active fragments of such peptides.

33. The method of claim 18 wherein the amphiphilic oligomer is selected from the group consisting of:

5



wherein  $n=3$  to 25 and  $m=1$  to 6;



wherein  $n=3$  to 25 and  $m=1$  to 7;



10

wherein  $n=3$  to 25,  $m=1$  to 7 and  $X=\text{O}$  or  $\text{N}$ ;




wherein  $m=0$  to 5 and  $\text{R}=\text{cholesterol}$  or  $\text{adamantane}$ ; or



wherein  $m=0$  to 5;

15



wherein  $m=0$  to 7;



wherein  $m=1$  to 7 and  $X=\text{N}$  or  $\text{O}$ .

Sub  
225  
cont

34. The method of claim 18 wherein the hydrophilic moiety is coupled to the hydrophobic moiety by a hydrolyzable bond.
35. The method of claim 18 wherein the hydrophilic moiety is coupled to the hydrophobic moiety by a non-hydrolyzable bond
- 5 36. A method for delivering a therapeutic compound across the blood-brain barrier comprising administering to a subject an amphiphilic drug-oligomer conjugate comprising a therapeutic compound conjugated to an oligomer, wherein the oligomer comprises a lipophilic moiety coupled to a hydrophilic moiety.
- 10 37. The method of claim 31 wherein the therapeutic compound is a peptide or protein.
- 15 38. The method of claim 31 wherein the therapeutic compound is a peptide and the peptide is selected from the group consisting of: enkephalin, adrenocorticotrophic hormone, adenosine deaminase ribonuclease, alkaline phosphatase, angiotensin, antibodies, arginase, arginine deaminase, asparaginase, caerulein, calcitonin, chemotrypsin, cholecystokinin, clotting factors, dynorphins, endorphins, endorphins, enkephalins, enkephalins, erythropoietin, gastrin-releasing peptide, glucagon, hemoglobin, hypothalamic releasing factors, interferon, katacalcin, motilin, neuropeptide Y, neurotensin, non-naturally occurring opioids, oxytosin, papain, parathyroid hormone, peptides prolactin, soluble CD-4, somatomedin, somatostatin, somatostatin, somatotropin, superoxide dismutase, thyroid stimulating hormone, tissue plasminogen activator, trypsin, vasopressin, and analogues and active fragments of such peptides.
- 20 39. The method of claim 31 wherein the therapeutic compound is [met<sup>5</sup>]enkephalin.
- 25 40. The method of claim 31 wherein the lipophilic moiety is coupled to the hydrophilic moiety by a hydrolyzable bond.

41. The method of claim 31 wherein the lipophilic moiety is coupled to the hydrophilic moiety by a non-hydrolyzable bond.

42. The method of claim 31 wherein the lipophilic moiety is coupled to the hydrophilic moiety by a bond selected from the group consisting of: amide bond, carbamate bond, carbonate bond and ester bond.

43. The method of claim 31 wherein the oligomer moiety is coupled to the drug moiety by a bond selected from the group consisting of amide bond, carbonate bond, carbamate bond, and ester bond.

44. The method of claim 31 wherein the lipophilic moiety is selected from the group consisting of fatty acids, C<sub>1-26</sub> alkyls, and cholesterol.

45. The method of claim 31 wherein the hydrophilic moiety is selected from the group consisting of sugars, and PEG<sub>1-7</sub>.

46. A method for inducing analgesia in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of an amphiphilic drug-oligomer conjugate comprising enkephalin conjugated to an oligomer, wherein the oligomer comprises a lipophilic moiety coupled to a hydrophilic moiety.

47. The method of claim 41 wherein the therapeutic compound is [met<sup>5</sup>]enkephalin.

48. The method of claim 41 wherein the lipophilic moiety is selected from the group consisting of fatty acids, C<sub>1-26</sub> alkyls, and cholesterol.

49. The method of claim 41 wherein the one or more hydrophilic moieties are selected from the group consisting of sugars and PEG.

50. The method of claim 41 wherein the hydrophilic moiety comprises a sugar and the sugar is selected from the group consisting of amino sugars and non-amino sugars.

51. The method of claim 41 wherein the oligomer is selected from the group consisting of:



wherein  $n=3$  to 25 and  $m=1$  to 6;




wherein  $n=3$  to 25 and  $m=1$  to 7;




wherein  $n=3$  to 25,  $m=1$  to 7 and  $X=\text{O}$  or  $\text{N}$ ;




wherein  $m=0$  to 5 and  $\text{R}=\text{cholesterol}$  or  $\text{adamantane}$ ; or



wherein  $m=0$  to 4 and  $\text{R}=\text{cholesterol}$  or  $\text{adamantane}$ ;



wherein  $m=0$  to 7;




wherein  $m=1$  to 7 and  $X=\text{N}$  or  $\text{O}$ .

6620T SECE460

sub  
part  
cont

52. An amphiphilic oligomer-enkephalin conjugate selected from the group consisting of:



53. A method for altering the binding affinity of a peptide to its receptor comprising  
 5 conjugating the peptide to an amphiphilic oligomer comprising a lipophilic moiety  
 coupled to a hydrophilic moiety.
54. The method according to claim 48 further characterized in that the binding affinity  
 is increased.
55. The method according to claim 48 further characterized in that the binding affinity  
 is reduced.
56. The method of claim 48 wherein the peptide is a peptide or protein.
57. The method of claim 48 wherein the peptide is selected from the group consisting  
 of: enkephalin, adrenocorticotrophic hormone, adenosine deaminase  
 15 ribonuclease, alkaline phosphatase, angiotensin, antibodies, arginase, arginine  
 deaminase, asparaginase, caerulein, calcitonin, chemotrypsin, cholecystokinin,  
 clotting factors, dynorphins, endorphins, endorphins, enkephalins, enkephalins,  
 erythropoietin, gastrin-releasing peptide, glucagon, hemoglobin, hypothalamic  
 20 releasing factors, interferon, kalacalcin, motilin, neuropeptide Y, neurotensin,  
 non-naturally occurring opioids, oxytocin, papain, parathyroid hormone, peptides  
 prolactin, soluble CD-4, somatomedin, somatostatin, somatostatin, somatotropin,  
 superoxide dismutase, thyroid stimulating hormone, tissue plasminogen  
 activator, trypsin, vasopressin, and analogues and fragments of such peptides.
58. The method of claim 48 wherein the peptides is [met<sup>5</sup>]enkephalin.
59. The method of claim 48 wherein the lipophilic moiety is selected from the group  
 25 consisting of fatty acids, C<sub>1-26</sub>alkyls, and cholesterol.
60. The method of claim 48 wherein the hydrophilic moiety is selected from the group  
 consisting of sugars or PEG<sub>1-7</sub>.

Sub  
 002  
 001

00420735-1090

61. The method of claim 48 wherein the oligomer is selected from the group consisting of:




wherein  $n=3$  to 25 and  $m=1$  to 6;



wherein  $n=3$  to 25 and  $m=1$  to 7;




wherein  $n=3$  to 25,  $m=1$  to 7 and  $X=\text{O}$  or  $\text{N}$ ;




10 wherein  $m=0$  to 5 and  $\text{R}=\text{cholesterol}$  or  $\text{adamantane}$ ; or



wherein  $m=0$  to 5;



wherein  $m=0$  to 7;



wherein  $m=1$  to 7 and  $X=\text{N}$  or  $\text{O}$ .

656207" SE/0E-160



62. The amphiphilic drug-oligomer conjugate of claim 1 wherein the therapeutic compound is a peptide and the peptide is selected from the group consisting of:

Ac-Phe-Arg-Trp-Trp-Tyr-Lys—NH<sub>2</sub> ;

5

Ac-Arg-Trp-Ile-Gly-Trp-Lys—NH<sub>2</sub> ;

Trp-Trp-Pro-Lys-His-Xaa—NH<sub>2</sub> ;

wherein Xaa is a naturally-occurring amino acid;

Trp-Trp-Pro-Xaa—NH<sub>2</sub> ,

wherein Xaa is Lys or Arg;

10

Tyr-Pro-Phe-Gly-Phe-Xaa—NH<sub>2</sub> ,

wherein Xaa is a naturally-occurring amino acid;

(D)Ile-(D)Met-(D)Ser-(D)Trp-(D)Trp-Gly<sub>n</sub>-Xaa—NH<sub>2</sub> ,

wherein n is 0 or 1 and wherein Xaa is Gly or the D-form of a naturally-occurring amino acid;

15

(D)Ile-(D)Met-(D)Thr-(D)Trp-Gly-Xaa—NH<sub>2</sub> ,

wherein Xaa is Gly or the D-form of a naturally-occurring amino acid;

Tyr-A1-B2-C3—NH<sub>2</sub> ,

wherein A1 is (D)Nve or (D)Nle,

sub  
aa7  
cont

B2 is Gly, Phe, or Trp, and

C3 is Trp or Nap;

Pm and red  $\{Me_xH_y-Tyr-(NMe)_z-Tyr-Xaa_z-NH_2\}$ ,

x is 0, 1, or 2,

y is 0, 1, or 2, and

z is 0 or 1, and

wherein Xaa is Phe, (D)Phe, or NHBzl, with the proviso that x and y together is never greater than 2;

Trp-Trp-Pro-D4-His<sub>z</sub>-Xaa<sub>z</sub>-NH<sub>2</sub>;

wherein z is 0 or 1,

wherein D4 is Lys or Arg, and

wherein Xaa is a naturally-occurring amino acid.

63. The method of claim 18 wherein the therapeutic compound is a peptide and the peptide is selected from the group consisting of:

Ac-Phe-Arg-Trp-Trp-Tyr-Lys-NH<sub>2</sub>;

Ac-Arg-Trp-Ile-Gly-Trp-Lys-NH<sub>2</sub>

Trp-Trp-Pro-Lys-His-Xaa-NH<sub>2</sub>

wherein Xaa is a naturally-occurring amino acid;

5

10

15

sub  
aa7  
cont

66207" SEZOE460

Trp-Trp-Pro-Xaa—NH<sub>2</sub>

wherein Xaa is Lys or Arg;

Tyr-Pro-Phe-Gly-Phe-Xaa—NH<sub>2</sub>

wherein Xaa is a naturally-occurring amino acid;

5

(D)Ile-(D)Met-(D)Ser-(D)Trp-(D)Trp-Gly<sub>n</sub>-Xaa—NH<sub>2</sub>

wherein n is 0 or 1 and wherein Xaa is Gly or the D-form of a naturally-occurring amino acid;

(D)Ile-(D)Met-(D)Thr-(D)Trp-Gly-Xaa—NH<sub>2</sub>

wherein Xaa is Gly or the D-form of a naturally-occurring amino acid;

10

Tyr-A1-B2-C3—NH<sub>2</sub>

wherein A1 is (D)Nve or (D)Nle,

B2 is Gly, Phe, or Trp, and

C3 is Trp or Nap;

15

Pm and red {Me<sub>x</sub>H<sub>y</sub>-Tyr-(NMe)<sub>z</sub>-Tyr-Xaa<sub>z</sub>—NH<sub>2</sub>},

x is 0, 1, or 2,

y is 0, 1, or 2, and

z is 0 or 1, and

sub  
aa7  
cont

wherein Xaa is Phe, (D)Phe, or NHBzl, with the proviso that x and y together is never greater than 2;

Trp-Trp-Pro-D4-His<sub>z</sub>-Xaa<sub>z</sub>-NH<sub>2</sub>,

wherein z is 0 or 1,

wherein D4 is Lys or Arg, and

wherein Xaa is a naturally-occurring amino acid.

sub  
aa  
aa  
aa

5

add  
aa  
aa  
aa

add  
c3

666207" SE 20E 450